

$B\bar{B}$ Mixing and CP Violation in $SU(2)_L \times SU(2)_R \times U(1)$ Models

Soo-hyeon Nam

Theoretical Physics Institute, University of Minnesota
Minneapolis, MN 55455

Abstract

We reexamine the mass mixing and CP violation in the $B\bar{B}$ system in general $SU(2)_L \times SU(2)_R \times U(1)$ models related to the recent measurements. The right-handed contributions can be sizable in $B\bar{B}$ mixing and CP asymmetry in B decays for a heavy W' even with a mass about 3 TeV. On the other hand the lower bound on the mass of W' can be taken down to approximately 300 GeV.

1 Introduction

The Standard $SU(2)_L \times U(1)$ model (SM) has been very successful in describing the known weak interaction phenomena. But the consistency of the present experimental results with the general scheme of charged weak interactions and CP violation in the SM is non-trivial so the model is challenged both experimentally and theoretically in its prediction of large CP violation effects in the B meson system [1]. As one of the simplest extensions of the Standard model gauge group and so the complement of the purely left-handed nature of the SM , the left-right theory with the group $SU(2)_L \times SU(2)_R \times U(1)$ has been widely studied. In this model, even with two generations of quarks one could get CP violation. With three generations of quarks, this model contains many parameters and many sources of CP violation [2]. One of the main source is the relative phase α between the two vacuum expectation values (VEVs) k and k' of the Higgs bidoublet Φ . The other sources are the complex phases in the left and right-handed quark mixing matrices U^L and U^R respectively. Here it would be convenient to regard U^L as the usual Cabbibo-Kobayashi-Maskawa (CKM) matrix and shift all phases except one to U^R . Using the Wolfenstein parameterization [3], we can express the CKM matrix approximately as

$$U^L = \begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4), \quad (1)$$

where λ (≈ 0.22) is an real expansion parameter, and A , ρ and η are also real quantities. From the above expression, the elements U_{ub}^L and U_{td}^L can be parameterized in terms of two phases γ and β respectively which form a unitary triangle (Fig.1) given by the orthogonality condition $\sum_{i=u,c,t} U_{id}U_{ib}^* = 0$. The recent significant measurements of β_{exp} give [4]

$$\sin 2\beta_{exp} = \begin{cases} 0.59 \pm 0.14 \pm 0.05 & (BABAR) \\ 0.99 \pm 0.14 \pm 0.06 & (Belle) \end{cases}. \quad (2)$$

If there are new physics effects involved, the experimental value β_{exp} can be expressed through other parameters representing the new physics as well as the phase of U_{td}^L in the SM .

In addition to the phases mentioned above, the masses (M_{W_R}) of the right-handed gauge bosons, the mixing angle ξ between the left- and right-handed gauge bosons W_L and W_R , and the right-handed gauge coupling constant g_R play an important role in new physics effects as fundamental input parameters in the left-right model (LRM). The success of the SM in the low-energy phenomenology requires that the masses (M_{W_R}) of the right-handed




Figure 1: Unitary triangle ($\lambda_i = U_{id}^* U_{ib}$).

gauge bosons are significantly larger than those (M_{W_L}) of left-handed gauge bosons. The first lower bound on M_{W_R} comes from a study of the low energy charged current sector allowing $M_{W_R} \gtrsim 3M_{W_L} \approx 240$ GeV [5]. Soon after, many theoretical limits have been presented on M_{W_R} and ξ under various assumptions [6]. The recent experimental limits are obtained by DØ and CDF from the direct search of the decay channels of the extra gauge bosons $W'^+ \rightarrow \ell_R^+ \nu_R$. DØ found $M_{W'} > 720$ GeV for $m_{\nu_R} \ll M_{W'}$ or $M_{W'} > 650$ GeV for $m_{\nu_R} = M_{W'}/2$ [7]. CDF has a limit of $M_{W'} > 652$ GeV for $m_{\nu_R} \ll M_{W'}$ if ν_R is stable [8]. All of these limits were obtained assuming manifest ($U^R = U^L$) or pseudo-manifest ($U^R = U^{L*} K$) left-right symmetry ($g_L = g_R$), where K is a diagonal phase matrix [9]. In this paper, we will not impose the discrete left-right symmetry which can cause troubles in explaining the cosmological baryon asymmetry and may lead to cosmological domain-wall problems [10]. However we will also consider the possibility of the left-right symmetric case among other possibilities.

The main purpose of this paper is to investigate the CP violation in the $B^0 \bar{B}^0$ system in the LRM related to the recent experiments, since $B^0 \bar{B}^0$ mixing has recently been advocated as a very sensitive probe for the CP violation and the presence of right-handed current. The SM contribution to $K^0 \bar{K}^0$ mixing was previously computed for any internal quark mass by Inami and Lim [11]. The right-handed contribution in the LRM has been done first by Beall, Bander, and Soni assuming the discrete left-right symmetry [12] and again by many authors [13] under various assumptions. But we notice that the contributions of the mixing angle ξ to $B^0 \bar{B}^0$ mixing and CP asymmetry can be large due to the heaviness of the top quark mass and the possibility of the enhancement in the right-handed quark mixing matrix in the general LRM . After reviewing the structure of the LRM in Sec.2, we will discuss $B^0 \bar{B}^0$ mixing in Sec.3 and CP asymmetry in B^0 decay in Sec.4 in detail.

2 $SU(2)_L \times SU(2)_R \times U(1)$ models

We briefly review here some of the main features of the *LRM*, which are needed to obtain our results. As the simplest extension of the *SM*, the gauge group of the *LRM* breaks down to that of the *SM* and it finally cascades down to $U(1)_{EM}$. The covariant derivative for the fermions $f_{L,R}$ with respect to the gauge group of the *LRM* appears as

$$D^\mu f_{L,R} = \partial^\mu f_{L,R} + ig_{L,R} W_{L,R}^{\mu a} T_{L,R}^a f_{L,R} + ig_1 B^\mu S f_{L,R}. \quad (3)$$

The electric charge which is the unbroken $U(1)$ generator is given as

$$Q = T_L^3 + T_R^3 + S. \quad (4)$$

The quarks and leptons transform under the gauge group of the *LRM* (T_L, T_R, S) as

$$\begin{aligned} q'_L &= \begin{pmatrix} u' \\ d' \end{pmatrix}_L \sim (\frac{1}{2}, 0, \frac{1}{6}), & q'_R &= \begin{pmatrix} u' \\ d' \end{pmatrix}_R \sim (0, \frac{1}{2}, \frac{1}{6}), \\ l'_L &= \begin{pmatrix} \nu' \\ e' \end{pmatrix}_L \sim (\frac{1}{2}, 0, -\frac{1}{2}), & l'_R &= \begin{pmatrix} \nu' \\ e' \end{pmatrix}_R \sim (0, \frac{1}{2}, -\frac{1}{2}), \end{aligned} \quad (5)$$

where the primes indicate that the fermions are gauge rather than mass eigenstates.

In order to generate masses for the fermions and implement the symmetry breaking, we need to include scalar fields into our theory. The simplest choice is to introduce one Higgs multiplet and two doublets,

$$\Phi = \begin{pmatrix} \phi_1^0 & \phi_1^+ \\ \phi_2^- & \phi_2^0 \end{pmatrix} \sim (\frac{1}{2}, \frac{1}{2}^*, 0), \quad \chi_L = \begin{pmatrix} \chi^+ \\ \chi^0 \end{pmatrix}_L \sim (\frac{1}{2}, 0, \frac{1}{2}), \quad \chi_R = \begin{pmatrix} \chi^+ \\ \chi^0 \end{pmatrix}_R \sim (0, \frac{1}{2}, \frac{1}{2}), \quad (6)$$

which acquire the vacuum expectation values (VEVs)

$$\langle \Phi \rangle = \begin{pmatrix} k & 0 \\ 0 & k' \end{pmatrix}, \quad \langle \chi_L \rangle = \begin{pmatrix} 0 \\ v_L \end{pmatrix}, \quad \langle \chi_R \rangle = \begin{pmatrix} 0 \\ v_R \end{pmatrix}, \quad (7)$$

where k and k' are complex, and v_L and v_R are real. χ_R is needed to generate a large M_{W_R} if $v_R \gg |k|, |k'|, v_L$. But χ_L is not essential unless we impose an left-right symmetry. It is also possible to adopt other choice of Higgs such as Higgs triplets instead [14]. The Lagrangian for the scalar field is

$$L_{scalar} = Tr[(D^\mu \Phi)^\dagger D_\mu \Phi] + (D^\mu \chi_L)^\dagger D_\mu \chi_L + (D^\mu \chi_R)^\dagger D_\mu \chi_R - V(\Phi, \chi_L, \chi_R). \quad (8)$$

For the Higgs fields described above, the kinetic terms in the Lagrangian generate the charged W boson matrix

$$M_{W^\pm}^2 = \begin{pmatrix} g_L^2(v_L^2 + K^2)/2 & -g_L g_R k^* k' \\ -g_L g_R k k'^* & g_R^2(v_R^2 + K^2)/2 \end{pmatrix} \equiv \begin{pmatrix} M_{W_L}^2 & M_{W_{LR}}^2 e^{i\alpha} \\ M_{W_{LR}}^2 e^{-i\alpha} & M_{W_R}^2 \end{pmatrix}, \quad (9)$$

where $K^2 = |k|^2 + |k'|^2$ and α is the phase of $k^* k'$. After the mass matrix is diagonalized by a unitary transformation the eigenvalues can be expressed in terms of a mixing angle as

$$\begin{aligned} M_W^2 &= M_{W_L}^2 \cos^2 \xi + M_{W_R}^2 \sin^2 \xi + M_{W_{LR}}^2 \sin 2\xi, \\ M_{W'}^2 &= M_{W_L}^2 \sin^2 \xi + M_{W_R}^2 \cos^2 \xi - M_{W_{LR}}^2 \sin 2\xi. \end{aligned} \quad (10)$$

Thus the mass eigenstates are written as

$$\begin{pmatrix} W^+ \\ W'^+ \end{pmatrix} = \begin{pmatrix} \cos \xi & e^{-i\alpha} \sin \xi \\ -\sin \xi & e^{-i\alpha} \cos \xi \end{pmatrix} \begin{pmatrix} W_L^+ \\ W_R^+ \end{pmatrix}, \quad (11)$$

where ξ is a mixing angle defined by

$$\tan 2\xi = -\frac{2M_{W_{LR}}^2}{M_{W_R}^2 - M_{W_L}^2}. \quad (12)$$

For $v_R \gg |k|, |k'|, v_L$, the mass eigenvalues and the mixing angle reduce to

$$M_W^2 \approx \frac{1}{2} g_L^2 (v_L^2 + K^2), \quad M_{W'}^2 \approx \frac{1}{2} g_R^2 v_R^2, \quad \xi \approx \frac{2g_L |k^* k'|}{g_R v_R^2}. \quad (13)$$

Here, the Schwarz inequality requires that $\zeta \equiv M_W^2/M_{W'}^2 \geq \xi_g \equiv (g_L/g_R)\xi$. From the limits on deviations of muon decay parameters from the V-A prediction, the lower bound on $M_{W'}$ can be obtained as follows [15]

$$(g_R/g_L)^2 \zeta < 0.033 \quad \text{or} \quad M_{W'} > (g_R/g_L) \times 440 \text{ GeV}. \quad (14)$$

We will use this number for our numerical analysis.

As well as the above charged gauge bosons, the charged would-be Goldstone bosons corresponding to the longitudinal components of the physical bosons take part in the charged current interactions. The coupling of the Goldstone fields to the fermions can be found from the detailed structure of the Higgs potential $V(\Phi, \chi_L, \chi_R)$ and the Yukawa couplings. However one can directly determine the Goldstone couplings in terms of the gauge couplings without considering the Higgs potential, but using the Ward identities that ensure that the




Figure 2: Tree-level Feynman diagrams for the gauge-bosons (W, W') and the Goldstone bosons (φ, φ') exchange.

unphysical poles in the two diagrams shown in Fig.2 should cancel each other [16]. The charged interaction Lagrangian is then given by

$$\begin{aligned}
L_{CC} = & -\frac{1}{\sqrt{2}}\bar{P}\gamma^\mu \left\{ [U^L g_L c_\xi L + U^R g_R s_\xi^+ R] W_\mu^+ + [-U^L g_L s_\xi L + U^R g_R c_\xi^+ R] W_\mu'^+ \right. \\
& + [(U^L M_P g_L c_\xi - U^R M_N g_R s_\xi^+) L + (-U^L M_N g_L c_\xi + U^R M_P g_R s_\xi^+) R] \frac{\varphi_\mu^+}{M_W} \\
& + [-(U^L M_P g_L s_\xi + U^R M_N g_R c_\xi^+) L + (U^L M_N g_L s_\xi + U^R M_P g_R c_\xi^+) R] \frac{\varphi_\mu'^+}{M_{W'}} \left. \right\} N \\
& + h.c. + \dots,
\end{aligned} \tag{15}$$

where c_ξ (s_ξ) $\equiv \cos \xi$ ($\sin \xi$), $s_\xi^\pm \equiv e^{\pm i\alpha} \sin \xi$, $L, R \equiv (1 \mp \gamma^5)/2$ denote left- and right-handed projection operators, $M_P = \text{diag}(m_u, m_c, m_t)$ and $M_N = \text{diag}(m_d, m_s, m_b)$ are the diagonalized quark mass matrices, P (N) is the mass eigenstate corresponding to its eigenvalue M_P (M_N), and U^L (U^R) is the left (right)-handed quark mixing matrix.




Figure 3: Box diagrams for $B^0 \bar{B}^0$ mixing with the gauge-bosons (W, W') and the Goldstone bosons (φ, φ').

3 $B^0 \bar{B}^0$ mixing

The effective Hamiltonian in the $B^0 \bar{B}^0$ system is obtained by integrating out the internal loop in the box diagrams in Fig.3 just as in the SM . We neglect external momenta and

d -quark mass, but the result is valid for general internal quark masses. One finds, using the Feynman-'t Hooft gauge, the charged gauge boson and Goldstone boson contributions to $B^0\bar{B}^0$ mixing in a straightforward manner :

$$H_{eff}^{B\bar{B}} = H_{eff}^{SM} + H_{eff}^{RR} + H_{eff}^{LR} \quad (16)$$

with

$$H_{eff}^{SM} = \frac{G_F^2 M_W^2}{4\pi^2} \sum_{i,j=u,c,t} \lambda_i^{LL} \lambda_j^{LL} \left\{ \left[\left(1 + \frac{x_i^2 x_j^2}{4}\right) f(x_i^2, x_j^2; 1) - 2x_i^2 x_j^2 g(x_i^2, x_j^2; 1) \right] (\bar{d}_L \gamma_\mu b_L)^2 \right. \\ \left. + x_b^2 x_i^2 x_j^2 g(x_i^2, x_j^2; 1) (\bar{d}_L b_R)^2 \right\}, \quad (17)$$

$$H_{eff}^{RR} = \frac{G_F^2 M_W^2}{4\pi^2} \left(\frac{g_R}{g_L}\right)^4 \sum_{i,j=u,c,t} \lambda_i^{RR} \lambda_j^{RR} \zeta f(x_i^2 \zeta, x_j^2 \zeta; 1) (\bar{d}_L \gamma_\mu b_R)^2, \quad (18)$$

$$H_{eff}^{LR} = \frac{G_F^2 M_W^2}{2\pi^2} \left(\frac{g_R}{g_L}\right)^2 \sum_{i,j=u,c,t} \left\{ \lambda_i^{LR} \lambda_j^{RL} x_i x_j \zeta [4g(x_i^2, x_j^2; \zeta) - f(x_i^2, x_j^2; \zeta)] (\bar{d}_L b_R) (\bar{d}_R b_L) \right. \\ + \lambda_i^{LL} \lambda_j^{LR} x_j x_b \xi_g^+ [x_i^2 (g(x_i^2, x_j^2; 1) - \frac{1}{4} f(x_i^2, x_j^2; 1)) (\bar{d}_L \gamma_\mu b_L)^2 \\ + (f(x_i^2, x_j^2; 1) - x_i^2 x_j^2 g(x_i^2, x_j^2; 1)) (\bar{d}_L b_R)^2] \\ + \lambda_i^{LL} \lambda_j^{RL} x_j x_b \xi_g^- [x_i^2 (g(x_i^2, x_j^2; 1) - \frac{1}{4} f(x_i^2, x_j^2; 1)) (\bar{d}_L \gamma_\mu b_L) (\bar{d}_R \gamma_\mu b_R) \\ \left. + (f(x_i^2, x_j^2; 1) - x_i^2 x_j^2 g(x_i^2, x_j^2; 1)) (\bar{d}_L b_R) (\bar{d}_R b_L) \right] \Big\}, \quad (19)$$

where

$$\frac{G_F}{\sqrt{2}} \equiv \frac{g_L^2}{8M_W^2}, \quad \xi_g^\pm \equiv e^{\pm\alpha} \xi_g, \quad \lambda_i^{AB} \equiv U_{id}^{A*} U_{ib}^B, \quad x_i \equiv \frac{m_i}{M_W} \quad (i = u, c, t), \quad (20)$$

and

$$f(x_i, x_j; \zeta) = \frac{\ln(1/\zeta)}{(1-\zeta)(1-x_i\zeta)(1-x_j\zeta)} + \left(\frac{x_i^2 \ln x_i}{(x_i - x_j)(1-x_i)(1-x_i\zeta)} + (i \longrightarrow j) \right), \\ g(x_i, x_j; \zeta) = \frac{\zeta \ln(1/\zeta)}{(1-\zeta)(1-x_i\zeta)(1-x_j\zeta)} + \left(\frac{x_i \ln x_i}{(x_i - x_j)(1-x_i)(1-x_i\zeta)} + (i \longrightarrow j) \right). \quad (21)$$

Although the form of the charged interactions in eqs.(17-19) is independent of our particular choice of scalar representation, the Ward identities require that the box diagrams contributing to $B^0\bar{B}^0$ mixing in the LRM are not gauge invariant [17]. In order to impose gauge invariance into our theory, we need to involve flavor-changing neutral Higgs bosons, but it has been known that their contributions, even at the tree-level as long as the mass of the flavor-changing Higgs boson is much heavier than $M_{W'}$,¹ are suppressed by approximately

¹The tree-level flavor-changing neutral Higgs contributions with masses M_H of order $M_{W'}$ in the manifest or pseudo-manifest left-right symmetric model were discussed in Ref. [13].

a factor of ζ compared to the above gauge boson contributions [18]. Therefore the above results, in the approximation of neglecting external momenta and d -quark mass, provide the complete effective Hamiltonian contributing to $B^0\bar{B}^0$ mixing.

At this stage, in order to analyze the obtained effective Hamiltonian quantitatively, we need to consider specific forms of the right-handed quark mixing matrices U^R . If the model has manifest or pseudo-manifest left-right symmetry, W_R mass has a stringent bound $M_{W_R} \geq 1.6$ TeV [12], and the W_R boson contributions to $B^0\bar{B}^0$ mixing and tree level b decay are very small. But, in general, the form of U^R is not necessarily to be restricted to manifest or pseudo-manifest symmetric type, so the W_R mass limit can be lowered to approximately 300 GeV by taking the following forms of U^R [19];

$$U_I^R = \begin{pmatrix} e^{i\omega} & \sim 0 & \sim 0 \\ \sim 0 & c_R e^{i\alpha_1} & s_R e^{i\alpha_2} \\ \sim 0 & -s_R e^{i\alpha_3} & c_R e^{i\alpha_4} \end{pmatrix}, \quad U_{II}^R = \begin{pmatrix} \sim 0 & e^{i\omega} & \sim 0 \\ c_R e^{i\alpha_1} & \sim 0 & s_R e^{i\alpha_2} \\ -s_R e^{i\alpha_3} & \sim 0 & c_R e^{i\alpha_4} \end{pmatrix}, \quad (22)$$

where c_R (s_R) $\equiv \cos \theta_R$ ($\sin \theta_R$) ($0^\circ \leq \theta_R \leq 90^\circ$). Here the matrix elements indicated ~ 0 may be $\lesssim 10^{-2}$ and the unitarity requires $\alpha_1 + \alpha_4 = \alpha_2 + \alpha_3$. From the $b \rightarrow c$ semileptonic decays of the B mesons, we can get an approximate bound $\xi_g \sin \theta_R \lesssim 0.013$ by assuming $|U_{cb}^L| \approx 0.04$ [20].

The obtained effective Hamiltonians in eqs.(17-19) are then further simplified using the Glashow-Iliopoulos-Maiani (GIM) cancellation $\sum_{i=u,c,t} \lambda_i = 0$ and neglecting the u -quark mass :

$$H_{eff}^{SM} = \frac{G_F^2 M_W^2}{4\pi^2} (\lambda_t^{LL})^2 S(x_t^2) (\bar{d}_L \gamma_\mu b_L)^2, \quad (23)$$

$$H_{eff}^{LR} = \frac{G_F^2 M_W^2}{2\pi^2} \left(\frac{g_R}{g_L} \right)^2 \left\{ [\lambda_c^{LR} \lambda_t^{RL} x_c x_t \zeta A_1(x_t^2, \zeta) + \lambda_t^{LR} \lambda_t^{RL} x_t^2 \zeta A_2(x_t^2, \zeta)] (\bar{d}_L b_R) (\bar{d}_R b_L) \right. \\ \left. + \lambda_t^{LL} \lambda_t^{RL} x_b \xi_g^- [x_t^3 A_3(x_t^2) (\bar{d}_L \gamma_\mu b_L) (\bar{d}_R \gamma_\mu b_R) + x_t A_4(x_t^2) (\bar{d}_L b_R) (\bar{d}_R b_L)] \right\}, \quad (24)$$

where

$$S(x) = \frac{x(4 - 11x + x^2)}{4(1 - x)^2} - \frac{3x^3 \ln x}{2(1 - x)^3}, \\ A_1(x, \zeta) = \frac{(4 - x) \ln x}{(1 - x)(1 - x\zeta)} + \frac{(1 - 4\zeta) \ln \zeta}{(1 - \zeta)(1 - x\zeta)}, \\ A_2(x, \zeta) = \frac{4 - x}{(1 - x)(1 - x\zeta)} + \frac{(4 - 2x + x^2(1 - 3\zeta)) \ln x}{(1 - x)^2(1 - x\zeta)^2} + \frac{(1 - 4\zeta) \ln \zeta}{(1 - \zeta)(1 - x\zeta)^2}, \quad (25) \\ A_3(x) = \frac{7 - x}{4(1 - x)^2} + \frac{(2 + x) \ln x}{2(1 - x)^3},$$

$$A_4(x) = \frac{2x}{1-x} + \frac{x(1+x)\ln x}{(1-x)^2}.$$

Note that $S(x)$ is the usual Inami-Lim function, $A_1(x, \zeta)$ is obtained by taking the limit $x_c^2 = 0$, and H_{eff}^{RR} is suppressed because it is proportional to ζ^2 . Also, in the case of U_I^R , one can see that there is no significant contribution of H_{eff}^{LR} to $B^0 \bar{B}^0$ mixing, so we will concentrate on the second type U_{II}^R in this section.

The dispersive part of the $B^0 \bar{B}^0$ mixing matrix element can then be written as

$$M_{12} = M_{12}^{SM} + M_{12}^{LR} = M_{12}^{SM} \left\{ 1 + \left(\frac{g_R}{g_L} \right)^2 r_{LR} \right\}, \quad (26)$$

where

$$\left(\frac{g_R}{g_L} \right)^2 r_{LR} \equiv \frac{M_{12}^{LR}}{M_{12}^{SM}} = \frac{\langle \bar{B}^0 | H_{eff}^{LR} | B^0 \rangle}{\langle \bar{B}^0 | H_{eff}^{SM} | B^0 \rangle}. \quad (27)$$

For specific phenomenological estimates one needs the hadronic matrix elements of the operators in eq.(23,24) in order to evaluate the mixing matrix element. We use the following parametrization :

$$\begin{aligned} \langle \bar{B}^0 | (\bar{d}_L \gamma_\mu b_L)^2 | B^0 \rangle &= \frac{1}{3} B_1 f_B^2 m_B, \\ \langle \bar{B}^0 | (\bar{d}_L \gamma_\mu b_L)(\bar{d}_R \gamma_\mu b_R) | B^0 \rangle &= -\frac{5}{12} B_2 f_B^2 m_B, \\ \langle \bar{B}^0 | (\bar{d}_L b_R)(\bar{d}_R b_L) | B^0 \rangle &= \frac{7}{24} B_3 f_B^2 m_B, \end{aligned} \quad (28)$$

where

$$\langle 0 | \bar{d}_\beta \gamma^\mu b_\alpha | B^0 \rangle = - \langle \bar{B}^0 | \bar{d}_\beta \gamma^\mu b_\alpha | 0 \rangle = - \frac{i f_B p_B^\mu}{\sqrt{2} m_B} \frac{\delta_{\alpha\beta}}{3}, \quad (29)$$

and where f_B is the B meson decay constant and B_i ($i = 1, 2, 3$) are the bag factors. In the vacuum-insertion method [21], $B_i = 1$ in the limit $m_b \simeq m_B$. We will use $f_B B_i^{1/2} = (210 \pm 40)$ MeV for our numerical estimates [22]. Using the standard values of the quark masses and $|U_{cd}^L| \approx 0.222$, one can express r_{LR} in terms of the mixing angle and phases in (22) as

$$\begin{aligned} r_{LR} \approx & l \left\{ 18.1 l \left(\frac{1 - \zeta - (3.49 - 14.0\zeta) \ln(1/\zeta)}{1 - 5.68\zeta} \right) \zeta_{SR}^2 e^{i\delta_1} \right. \\ & \left. - 739 \left(\frac{1 - 5.04\zeta - (0.483 - 1.93\zeta) \ln(1/\zeta)}{1 - 10.4\zeta + 31.3\zeta^2} \right) \zeta_{SRCR} e^{i\delta_2} - 7.68 \xi_{gSR} e^{i\delta_3} \right\}, \end{aligned} \quad (30)$$

where $l = 0.009/|U_{td}^L|$, $\delta_1 = -2\beta + \alpha_2 - \alpha_3$, $\delta_2 = -\beta - \alpha_3 + \alpha_4$, $\delta_3 = -\beta - \alpha_3$ and the mixing phase α was absorbed in α_i by redefining $\alpha_i + \alpha \rightarrow \alpha_i$.




Figure 4: Behavior of the ratio $|r_{LR}|$ as $\delta_{1,2}$ are varied.




Figure 5: Allowed region for $|r_{LR}|$ and θ_R .

Now we investigate numerically the behavior of the ratio $|r_{LR}|$, which is the deviation of M_{12} from the SM , under variation of $M_{W'}$, ξ_g , θ_R and the phases in U^R , assuming $l = 1$. Although we use the average value of $|U_{td}|$ which might be different from the actual value of $|U_{td}^L|$, it should not affect the order of magnitude in our estimates. First, in order to see the dependence of $|r_{LR}|$ on the phases, we fix $M_{W'} = 800$ GeV, $\xi_g = 0.005$, $\theta_R = 15^\circ$, and set $\delta_3 = \pi$ because its effect is relatively much smaller than that of δ_1 and δ_2 . The plot is shown in Fig.4. From eq.(30) and Fig.4, one can see that $|r_{LR}|$ becomes maximal when $\delta_{1,3} = \pi$ and $\delta_2 = 0$, and minimal when $\delta_{1,2,3} = \pi$ if $\theta_R \lesssim 70^\circ$ (or $\delta_{1,2} = \pi$ and $\delta_3 = 0$ if $\theta_R \gtrsim 70^\circ$). This behavior also holds for other values of $M_{W'}$ and ξ_g . Since $|r_{LR}|$ is the continuously varying function of the phases, we can probe the allowed region for $|r_{LR}|$ with respect to

the parameters $M_{W'}$, ξ_g and θ_R . Next, we fix $M_{W'} = 800$ GeV, $\xi_g = 0.005$, and evaluate $|r_{LR}|$ by varying θ_R . Note that $|r_{LR}|$ can approach zero at a non-zero θ_R near 73° as shown in Fig.5. Otherwise, it is larger than 1, which means that generally it is possible to have $|M_{12}^{LR}| \gg |M_{12}^{SM}|$. In Fig.6, we consider the behavior of $|r_{LR}|$ for $g_R/g_L \geq 0.5$, $\xi_g = 0.0004$ and $\theta_R = 14^\circ, 70^\circ$ as $M_{W'}$ is varied. The behavior of $|r_{LR}|$ exhibits a substantial dependence on $M_{W'}$, and $|r_{LR}|$ can be larger than 1 even for $M_{W'} \sim 2$ TeV. Moreover, it can be noticed that $|r_{LR}|$ falls near $M_{W'} \sim 300$ GeV at certain angles and phases in the mixing matrices. This reflects a possibility of relatively light masses of W' compared to the previously known bound. We will return to this point in Sec.4. The dependence of $|r_{LR}|$ on ξ_g satisfying $\xi_g \sin \theta_R \lesssim 0.013$ at fixed $M_{W'} = 700$ GeV and $\theta_R = 14^\circ, 70^\circ$ is shown in Fig.7. As one can see, $|r_{LR}|$ can be enhanced up to 10% of the SM contribution for the given inputs. Although its effect is smaller than that of other parameters, it is not negligible and can be dominant in $|r_{LR}|$ if the first two ζ dependent terms in eq.(30) cancel each other.




Figure 6: Allowed regions for $|r_{LR}|$ and $M_{W'}$ for $g_R/g_L \geq 0.5$. The dotted lines correspond to the lower bounds on $M_{W'}$ in eq.(14) for the ratio $g_R/g_L = 1, 2, 3$ and 4 respectively.




Figure 7: Allowed regions for $|r_{LR}|$ and ξ_g for $M_{W'} = 700$ GeV.

As we mentioned previously, the average value of $|U_{td}|$ might be different from the actual

value of $|U_{td}^L|$, and there is also ambiguity from errors in $f_B B_i^{1/2}$. Therefore the mass mixing ΔM_B^{SM} can be either much larger or smaller than ΔM_B^{exp} . However, if we assume that $0.5 \lesssim |\Delta M_B^{SM}/\Delta M_B^{exp}| \lesssim 2$, we can get specific bounds on the mass $M_{W'}$ and the angle θ_R using the experimental value $\Delta M_B^{exp} \simeq 0.472 \times 10^{12} s^{-1}$. We will estimate the lowest possible bound on $M_{W'}$ with respect to θ_R in their parameter space with the numerical consideration of $\sin 2\beta$ in the next section.

4 CP asymmetry in B^0 decay

The CP angle β in CKM matrix can be measured in $B \rightarrow J/\psi K_S$ decays. In B decays into a final CP eigenstate $J/\psi K_S$, β is related to a parametrization invariant quantity λ as follows [1]:

$$\sin 2\beta_{eff} = \text{Im}\lambda(B^0 \rightarrow J/\psi K_S), \quad (31)$$

where

$$\lambda \equiv -\left(\frac{q}{p}\right)_B \frac{A(\bar{B}^0 \rightarrow J/\psi K_S)}{A(B^0 \rightarrow J/\psi K_S)}, \quad \left(\frac{q}{p}\right)_B \simeq \frac{M_{12}^*}{|M_{12}|}. \quad (32)$$

The minus sign in the above expression is coming from the fact that $J/\psi K_S$ is CP -odd. As mentioned earlier, $\beta_{eff} = \beta$ in the SM .

In the LRM , the two types of U^R give us two distinct results. In the case of U_I^R , the W' contribution to the mixing parameter $(q/p)_B$ is negligible so that $(q/p)_B \simeq (q/p)_{SM} = e^{-2i\beta}$. Then the CP angle β_{eff} can be expressed by

$$\begin{aligned} \sin 2\beta_{eff}^I &\simeq -\text{Im} \left(e^{-2i\beta} \frac{U_{cs}^{L*} U_{cb}^L + (g_R/g_L)^2 (-2U_{cs}^{L*} U_{cb}^R \xi_g^+ + U_{cs}^{R*} U_{cb}^R \zeta)}{U_{cs}^L U_{cb}^{L*} + (g_R/g_L)^2 (-2U_{cs}^L U_{cb}^{R*} \xi_g^- + U_{cs}^R U_{cb}^{R*} \zeta)} \right) \\ &\simeq -\text{Im} \left(e^{-2i\beta} \frac{1 + 25(g_R/g_L)^2 (-2s_R \xi_g e^{i\alpha_2} + c_R s_R \zeta e^{i(\alpha_2 - \alpha_1)})}{1 + 25(g_R/g_L)^2 (-2s_R \xi_g e^{-i\alpha_2} + c_R s_R \zeta e^{-i(\alpha_2 - \alpha_1)})} \right), \end{aligned} \quad (33)$$

where the mixing angle α is absorbed in α_i again, and we ignored the $K\bar{K}$ mixing and assumed that

$$\begin{aligned} \langle J/\psi K_S | \bar{c}_L \gamma_\mu s_L \bar{b}_L \gamma^\mu c_L | B^0 \rangle &\simeq \langle J/\psi K_S | \bar{c}_R \gamma_\mu s_R \bar{b}_R \gamma^\mu c_R | B^0 \rangle \\ &\simeq -\frac{1}{2} \langle J/\psi K_S | \bar{c}_L \gamma_\mu s_L \bar{b}_R \gamma^\mu c_R | B^0 \rangle. \end{aligned} \quad (34)$$

As one can easily see in eq.(33), $\sin 2\beta_{eff} = \sin 2\beta$ if $\alpha_{1,2} = 0$ or π .

For illustration of the possible effect of the new interaction on the effective value of $\sin 2\beta$: $\sin 2\beta_{eff}$, we assume that the SM contribution produces $\sin 2\beta = 0.60$, and show the region




Figure 8: Behavior of $\sin 2\beta_{eff}$ as $\alpha_{1,2}$ are varied.




Figure 9: Contour plots corresponding to $\sin 2\beta_{eff} = 0.99$ for $\sin 2\beta = 0.60$, (a) $\xi_g = \zeta/2$ and $g_R = g_L$, (b) $\xi_g = \zeta$ and $g_R = g_L$, and (c) $\xi_g = \zeta/2$ and $g_R = 2g_L$. The dotted line corresponds to the lower bound on $M_{W'}$ for $g_R = 2g_L$.

of parameters, where the effective value is shifted to $\sin 2\beta_{eff} \sim 1$. We first plot $\sin 2\beta_{eff}$ in Fig.8 for the typical values $M_{W'} = 800$ GeV, $\xi_g = 0.005$, $\theta_R = 15^\circ$ and $g_R = g_L$ as $\alpha_{1,2}$ are varied. In the figure, $\sin 2\beta_{eff}$ has a maximum variation from $\sin 2\beta$ near $\alpha_2 \approx \pi/2$ and $\alpha_1 = \pi$, and this behavior holds for other values of $M_{W'}$, ξ_g and θ_R . Next, we plot the




Figure 10: Contour plots of ΔM_B (dashed) for $|U_{td}^L| = 0.009$ and those corresponding to $\sin 2\beta_{eff} = 0.99$ (dotted) for $\sin 2\beta = 0.60$.

contour corresponding to $\sin 2\beta_{eff} = 0.99$ satisfying $\xi_g \sin \theta_R \lesssim 0.013$ in the parameter space of $M_{W'}$ and θ_R for $\alpha_2 = \pi/2$, $\alpha_1 = \pi$, $\xi_g = \zeta/2, \zeta$ and $g_R/g_L = 1, 2$ in Fig.9.² As one can see, the upper bound of $M_{W'}$ goes down with decreasing ξ_g and g_R/g_L . Therefore, under the given assumption, $g_R \ll g_L$ is disfavored, and so is $M_{W'} \gg 1$ TeV unless $g_R \gg g_L$.

In the case of U_{II}^R , one has $U_{cs}^R \sim 0$ so that the ζ ($M_{W'}$) dependent term in eq.(33) is very small. However the ratio $(q/p)_B$ depends on $M_{W'}$. Thus the W' contribution enters in a somewhat different way.

$$\sin 2\beta_{eff}^{II} \simeq -\text{Im} \left(e^{-2i\beta} \frac{(1 + (g_R/g_L)^2 r_{LR}^*)}{|1 + (g_R/g_L)^2 r_{LR}|} \frac{(1 - 50(g_R/g_L)^2 s_R \xi_g e^{i\alpha_2})}{(1 - 50(g_R/g_L)^2 s_R \xi_g e^{-i\alpha_2})} \right). \quad (35)$$

Unlike in the previous case, we need to consider here the mass mixing of $B^0 \bar{B}^0$ in order to analyze $\sin 2\beta_{eff}$ numerically. Assuming that $|U_{td}^L| = 0.009$ and $\sin 2\beta = 0.60$, we plot the contours corresponding to $\sin 2\beta_{eff} = 0.99$ and $\Delta M_B^{LR} = \Delta M_B^{exp}$ in the parameter space of

²We did the same analysis for $g_R/g_L = 0.5$ but there was no allowed region.

$\delta_{1,2}$ for $\delta_3 = \pi$, $\xi_g = \zeta/4$ and $g_R/g_L = 0.5, 1, 2$ by varying θ_R and $M_{W'}$ in Fig.10. Because of the non-triviality of the behavior of $\sin 2\beta_{eff}$ on δ_i , we repeated this analysis until the two contours have overlapped by varying $M_{W'}$ from 350 GeV to 8 TeV, and found that it appeared where $350 \text{ GeV} \lesssim M_{W'} \lesssim 1.3 \text{ TeV}$ if $g_R/g_L = 0.5$, $440 \text{ GeV} \lesssim M_{W'} \lesssim 3.1 \text{ TeV}$ if $g_R/g_L = 1$ and $880 \text{ GeV} \lesssim M_{W'} \lesssim 7.1 \text{ TeV}$ if $g_R/g_L = 2$. Even though the existence of a heavy W' with the mass $M_{W'} > 7 \text{ TeV}$ may be allowed by the numerical analysis of ΔM_B^{LR} , it is excluded by that of $\sin 2\beta_{eff}$ under the given assumption. For the different values of ξ_g , we also have similar results.

5 Conclusion

In the *LRM*, if one does not impose the manifest or pseudo-manifest left-right symmetry, the W' contributions to $B^0\bar{B}^0$ mixing and CP asymmetry in B^0 decays are highly dependent upon the phases in the mass mixing matrix $U^{L,R}$. For certain phases, the contribution of W' with a heavy mass about a few TeV to $B^0\bar{B}^0$ mixing can be sizeable. On the other hand, there is also a possibility of the existence of W' with a light mass about a few hundred GeV, whose contribution can be either very large or small, and so the contribution of the mixing angle ξ is not negligible. Since the existence of a light W' requires a small g_R , $g_R \lesssim g_L$, one can see from eq.(26) and Fig.6 that its contribution is limited. Therefore even assuming that $\Delta M_B^{LR} \lesssim \Delta M_B^{SM}$, we find that there is a possibility of a light W' with a mass $M_{W'} \sim 300 \text{ GeV}$.

Also this possibility arises from the numerical analysis of the CP asymmetry in B^0 decay. Since U_{td}^L is not known with sufficient accuracy, estimates of the pure right-handed current contributions to ΔM_B and $\sin 2\beta$ are somewhat uncertain. But, for the certain values of the parameter sets, one can see from Fig.9 and Fig.10 that the CP asymmetry parameter $\sin 2\beta$ can be as large as almost 1, and the mass of W' can be as small as about 350 GeV. Therefore, the existence of the light W' can be tested once future experiments confirm the value of $\sin 2\beta$ and $|U_{td}^L|$.

6 Acknowledgements

The author would like to thank M. B. Voloshin for helpful comments.

References

- [1] For a recent review, see I. I. Bigi and A. I. Sanda, *CP Violation* (Cambridge University, Cambridge, UK, 2000).
- [2] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974); R. N. Mohapatra and J. C. Pati, *ibid.* **11**, 566 (1975); **11**, 2558 (1975).
- [3] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
- [4] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 091801 (2001); Belle Collaboration, K. Abe *et al.*, *ibid.* **87**, 091801 (2001).
- [5] M. A. B. Bég, R. V. Budny, R. N. Mohapatra, and A. Sirlin, Phys. Rev. Lett. **38**, 1252 (1977).
- [6] For a review, see P. Langacker and S. U. Sankar, Phys. Rev. D **40**, 1569 (1989).
- [7] DØ Collaboration, S. Abachi *et al.*, Phys. Rev. Lett. **76**, 3271 (1996).
- [8] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **74**, 2900 (1995).
- [9] For a review, see R. N. Mohapatra, *Unification and Supersymmetry* (Springer, New York, 1992).
- [10] P. Langacker, in *CP violation*, edited by C. Jarlskog (World Scientific, Singapore, 1989).
- [11] T. Inami and C. S. Lim, Prog. Theor. Phys. **65**, 297 (1981).
- [12] G. Beall, M. Bander, and A. Soni, Phys. Rev. Lett. **48**, 848 (1982).
- [13] R. N. Mohapatra, G. Senjanović, and M. D. Tran, Phys. Rev. D **28**, 546 (1983); R. Decker and U. Türke, Z. Phys. C **26**, 117 (1984); G. Ecker and W. Grimus, *ibid.* **30**, 293 (1986); P. Ball, J.-M. Frère, and J. Matias, Nucl. Phys. **B572**, 3 (2000).
- [14] R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980); Phys. Rev. D **23**, 165 (1981).
- [15] B. Balke *et al.*, Phys. Rev. D **37**, 587 (1988).
- [16] K. Fujikawa, B. W. Lee, and A. I. Sanda, Phys. Rev. D **6**, 2923 (1972).

- [17] D. Chang, J. Basecq, L.-F. Li, and P. B. Pal, Phys. Rev. D **30**, 1601 (1984).
- [18] W.-S. Hou and A. Soni, Phys. Rev. D **32**, 163 (1985); J. Basecq, L.-F. Li, and P. B. Pal, *ibid.* **32**, 175 (1985).
- [19] F. I. Olness and M. E. Ebel, Phys. Rev. D **30**, 1034 (1984); D. London and D. Wyler, Phys. Lett. B **232**, 503 (1989); also see Ref. [6].
- [20] M. B. Voloshin, Mod. Phys. Lett. A **12**, 1823 (1997).
- [21] M. K. Gaillard and B. W. Lee, Phys. Rev. D **10**, 897 (1974).
- [22] C. W. Bernard *et al.*, Phys. Rev. Lett. **81**, 4812 (1998).